

Chasing electric flux in hot QCD

Y. Schröder

Faculty of Physics, University of Bielefeld, D-33501 Bielefeld, Germany

Abstract

In this talk, I present the status of attempts to analyze the behavior of the so-called spatial 't Hooft loop, which can be taken as an order parameter for the deconfinement phase transition in pure $SU(N)$ gauge theory. While lattice data show a strikingly universal scaling of extracted k -string tensions for various values of k and N , the analytic approach to these observables might need some refinement.

Key words: Thermal field theory, Perturbative QCD, Quark-gluon plasma
PACS: 11.10.Wx, 12.38.Bx, 12.28.Mh

1. Introduction

Four-dimensional $SU(N)$ gauge theory undergoes a phase transition from a cold and confined phase into a hot and deconfined phase at a critical temperature T_c , which is intimately related to the breaking of the $Z(N)$ centergroup symmetry. In this respect, a widely used order parameter is the Polyakov loop $P = \frac{1}{N} \text{Tr} \mathcal{P} \exp(ig \int_0^{1/T} dx_0 A_0)$, whose thermal average $\langle P \rangle \neq 0$ in the deconfined phase. There are however some formal issues associated with the use of $\langle P \rangle$ as order parameter: P cannot be defined at strictly zero temperature, and its lattice-discretized version exhibits ultraviolet divergences in the continuum limit.

Analyzing other choices, and noting that the spatial Wilson loop shows only area law behavior at $T \neq 0$, while at $T = 0$ it exhibits area/perimeter law behavior for broken/unbroken $Z(N)$ symmetry, it has been proposed to use the 't Hooft flux loop operator [1] – which in some sense is dual to Wilson loops – as an alternative order parameter [2].

In order to give a definition of the 't Hooft flux loop operator, consider measuring color-electric flux through a large surface S , in $SU(N)$ gauge theory at finite temperature. Since the color-electric field $(\mathbf{E})_i = F_{0i}$ is in the Lie Algebra of $SU(N)$, one asks for its flux projected onto some special direction Y in $SU(N)$: $\Phi = \frac{1}{g} \int_S d\mathbf{S} \cdot \text{Tr} \mathbf{E} Y$. Note that




Fig. 1. Comparison of lattice data and analytical results for ratios of the k -string tensions of Eq. (2) as a function of temperature, for different gauge groups. The horizontal lines are analytic leading-order (LO) results $\frac{k(N-k)}{(N-1)}$, the curved lines are NNLO with $T_c/\Lambda_{\overline{\text{MS}}} = (1.1, 1.35)$. The plots are from ref. [4].

$\langle \Phi \rangle = 0$. Now define the 't Hooft loop (or E-flux) operator [1,2] $V = \exp(4\pi i \Phi)$, whose thermal average $\langle V \rangle$ behaves non-trivially.

The directions (in matrix space) Y are constrained by gauge invariance. Since $F_{\mu\nu}^\Omega = \Omega F_{\mu\nu} \Omega^{-1}$ transforms as an adjoint, one has to choose special Y 's, namely generators of the center of $SU(N)$ which respect $\exp(2\pi i Y_k) = \exp(2\pi i \frac{k}{N}) \mathbb{1}_{N \times N}$ for $k = 0, \dots, N-1$. These are not unique; one simple choice [3] is $Y_k = \frac{1}{N} \text{diag}(\{k\}^{N-k}, \{k-N\}^k)$. One hence obtains a set of gauge invariant (for a detailed proof¹ see [7]) operators

$$V_k = \exp \left(\frac{4\pi i}{g} \int_S d\mathbf{S} \cdot \text{Tr} \mathbf{E} Y_k \right). \quad (1)$$

One can finally define the thermal averages

$$\langle V_k \rangle \sim \exp(-\sigma_k(T) A_S), \quad (2)$$

which at large areas A_S of the surface S define the k -string tensions σ_k .

The tensions σ_k can be computed either with analytic methods in a weak-coupling expansion (see Sec. (2)), or on the lattice [4]. From Fig. (1), one observes fairly good agreement between analytic and numerical results for the dimensionless ratios σ_k/σ_1 .

Interestingly, when choosing a different normalization on the vertical axis, on the lattice one clearly observes more violation of “Casimir scaling” at low temperature [4], as shown in Fig. (2). Moreover, and most strikingly, the lattice data for different gauge groups and

¹ To prove that V_k acting on physical states is gauge invariant, the main idea is to construct the physical subspace spanned by gauge invariant combinations of gauge fields (which are Wilson loops $W = \text{Tr} \mathcal{P} \exp(i \oint_C d\mathbf{l} \cdot \mathbf{A})$), to show $V_k W V_k^\dagger = \exp(2\pi i Y_k) W$ and therefore $V_k^\Omega W (V_k^\Omega)^\dagger = \exp(2\pi i \Omega Y_k \Omega^{-1}) W = \Omega \exp(2\pi i Y_k) \Omega^{-1} W = \exp(2\pi i Y_k) W$, such that when acting on a physical state constructed from one Wilson loop one obtains $V_k^\Omega | \text{phys} \rangle = V_k^\Omega W (V_k^\Omega)^\dagger V_k^\Omega | 0 \rangle$.




Fig. 2. The same data as in Fig. (1), with a different normalization. Note the universality for different N , k . Solid line: NLO σ_k with 2-loop running coupling, at $T_c/\Lambda_{\overline{MS}} = 1.35$. The plot is taken from ref. [4].

different values of k appear to fall onto a universal curve, hinting towards (large- N ?) universality. Clearly, it would be nice to understand this behavior analytically.

2. Analytic results

The weak-coupling expansion has been driven to next-to-next-to leading order (NNLO),

$$\frac{\sigma_k}{T^2} = \frac{4\pi^2}{3\sqrt{3}g^2N} k(N-k) \left(1 - 15.2785 \dots \frac{g^2N}{(4\pi)^2} + c_3(N,k) \frac{g^3N^{\frac{3}{2}}}{(4\pi)^3} + c_4(N,k) \frac{g^4N^2}{(4\pi)^4} \right). \quad (3)$$

At leading order, one observes Casimir scaling, while the NLO coefficient [5] scales in the same way with N and k , provided the expansion is organized in terms of g^2N .

The NNLO coefficient $c_3(N,k)$ has been determined in [6]. It is small (which is the reason why, for simplicity, in Fig. (2) only NLO is shown); it contributes (in analogy to e.g. weak-coupling results for the QCD pressure [8]) with a sign opposite to the g^2 term; its nonanalytic (in α_s) magnitude g^3 is well-understood, originating from infrared sensitive $\mathcal{O}(g^4)$ graphs that necessitate an all-order resummation of this specific sector; and it is expressed in [6] as a specific two-dimensional integral plus an infinite sum.

From the analytic side, the question now clearly is whether the unknown NNNLO term $c_4(N,k)$ can be computed, and whether it will help in explaining the apparent discrepancy at low temperatures $T \leq 4T_c$ between lattice and analytic results as shown in Fig. (2). To set our minds for doing this, it might be useful to recall the setup in which the analytic results that are known so far have been obtained.

The basic idea is to compute σ_k as tunneling effect through a perturbatively calculable potential barrier. Hence, one first needs to compute an effective potential. The presence of V_k breaks the center group symmetry, since it chooses specific directions Y_k . This leads to the Polyakov loop P picking up a phase $\exp(2\pi i \frac{k}{N})$ when moving through the surface S (at, say, $z = 0$, if the S is simply chosen as spanned by an $L_x \times L_y$ loop in the xy plane), which means that the field component $A_0(z)$ is discontinuous at S . One

can parameterize this behavior with (diagonal, traceless, $N \times N$) matrices C , according to $P \equiv \text{Tr} \exp(iC(z))$. Now, constraining the path integral to these allowed values of A_0 with a delta function one obtains

$$\langle V_k \rangle = \int [\mathcal{D}A_0][\mathcal{D}\mathbf{A}] \delta(P - \text{Tr} \exp(iC(z))) \exp(-S[A]) \sim \exp(-L_x L_y \sigma_k),$$

where the last step is valid in the limit of large loops and defines the minimum value of the effective potential U , with $\sigma_k = \min_{\{C(z)\}} U[C(z)]$. This minimization is taken with respect to all $C(z)$ which have a $2\pi k$ discontinuity at $z = 0$ and vanish as $z \rightarrow \infty$.

The minimal profile of the effective potential U is conjectured to be realized along the simplest path in the space of the matrices $C \sim 2\pi q Y_k$, with $0 < q < 1$ (while there is no general proof yet, this has been shown at large N and for $N = 3, 4$). To compute U in a perturbative expansion around the Polyakov loop constraint, one sets $A_0 = C(z) + gQ_0$ and $A_i = gQ_i$, chooses background field gauge $S_{\text{gf}} = \frac{1}{\xi} \text{Tr}[D_\mu(C)Q_\mu]^2$, performs the loop expansion, and finally minimizes with respect to q . On the technical side, the q -dependence represents a smooth interpolation between bosonic and fermionic Matsubara sums [9]. Additionally, going beyond NLO entails resumming the IR enhanced g^4 diagrams, which is most transparently done in the framework of dimensional reduction. As the magnetic sector does not contribute before g^5 , the relative order g^4 is computable.

3. Conclusions

The spatial 't Hooft loop V_k measures the color-electric flux, which at large temperatures is due to free screened gluons. This induces an area law behavior, whose corresponding tensions $\sigma_k \sim \ln \langle V_k \rangle$ are hard to measure on the lattice, since they are exponentially suppressed, and hence trigger a sampling problem that increases with the loop size. Existing lattice measurements indicate universal behavior of the σ_k , while NNLO weak-coupling results seem to correctly predict ratios of σ_k , but not their normalization close to T_c . Hope is expressed that knowledge of the NNNLO term would cure this lack of predictivity, and the setup in which to obtain this contribution is roughly sketched. As a final remark, let us stress that the ratios σ_k/σ_1 can serve as stringent tests for formulations of Yang–Mills theory derived from e.g. string theory.

References

- [1] G. 't Hooft, Nucl. Phys. B **138** (1978) 1.
- [2] A. Kovner and B. Rosenstein, Int. J. Mod. Phys. A **7** (1992) 7419.
- [3] C. Korthals Altes and A. Kovner, Phys. Rev. D **62** (2000) 096008.
- [4] P. de Forcrand, B. Lucini and D. Noth, PoS **LAT2005** (2006) 323.
- [5] T. Bhattacharya, A. Gocksch, C. Korthals Altes and R. D. Pisarski, Phys. Rev. Lett. **66** (1991) 998; Nucl. Phys. B **383** (1992) 497.
- [6] P. Giovannangeli and C. Korthals Altes, Nucl. Phys. B **721** (2005) 1; Nucl. Phys. B **721** (2005) 25.
- [7] C. Korthals Altes, lecture notes, Bielefeld Graduate School, Sept 2007; see <http://www.physik.uni-bielefeld.de/igs/schools/Fall12007/altes.writeup.pdf>.
- [8] J. I. Kapusta, Nucl. Phys. B **148** (1979) 461.
- [9] Y. Schröder, arXiv:0807.0500 [hep-ph].